

## Effect of different levels of buds load on bud behavior and fruit quality of Early Sweet grapevine

Thoraua S. A. Abo-ELwafa  
Viticulture. Dept. Hort. Res. Institute. Agric. Res. Center, Giza, Egypt  
Corresponding author: thorayaaboelwafa@yahoo.com

### Abstract

This investigation was carried out during both 2016 & 2017 seasons in a private vineyard at El-Khatatba region, Minufya Governorate to study the effect of different levels of buds load on bud behavior and the quality of Early Sweet grapes. The chosen vines were 7-year-old, grown in a sandy soil, spaced at 2 X 3 meters, irrigated by the drip irrigation system, trellised by the Spanish Parron shape system. Vines were trained to quadrilateral cordon and spur-pruned. Vines pruned to three different levels of bud load (48, 72 and 96 buds/vine) and number of buds per fruiting spurs adopted to 2, 4 and 6 buds/spur. The results showed that (48 buds/vine) 24 spurs  $\times$  2 buds increased bud behavior, physical characteristics, vegetative growth, wood ripening and weight of pruning. In addition, total carbohydrates in canes, however, the level load of (96 buds/vine) 24 spurs  $\times$  4 buds gave the highest yield per vine in both seasons. Finally, the best results was (48 buds/vine) namely 24, 12 and 8 fruiting spurs adopted to 2, 4 and 6 buds/spur, respectively.

**Keywords:** Grapevines, Early Sweet, spur pruned, vegetative growth, yield and quality

### Introduction

Grape is considered one of the most important fruit crops all over the world and popular and favorite fruit crops. Moreover, it is considered to be the second most important fruit crop after citrus in Egypt as its acreage, production and exportation in Egypt.

The production of grapes in Egypt increased as a new varieties became known and culture practices post-harvest handling as well as a new marketing methods utilized in the new reclaimed areas in the Egyptian deserts particularly the early ripening cultivars such as "Flame seedless", "Superior" and "Early Sweet

Early sweet grapevine cv. is considered a prime and outstanding grapevine cv. grown under Egypt conditions. It ripens early especially when treated with breakages and easily marketing to most foreign countries. Early Sweet Grapes is a large seedless berry with a creamy white color. The high sugar level gives this grape a sweet flavor with a hint of Muscat and an extremely high juice level.

Pruning is the most important cultural practice in the management of grapevine to sustain production and productivity. Pruning methods have been developed to balance fruit productivity, vegetative growth and attain maximum yield without reducing vine vigour. An increase in the severity of pruning will increase the vigor of individual shoot at the expense of total growth and crop (Weaver, 1976 and Celiket *et al.*, 1998). Pruning practice through which grape production can be increased and cluster quality improved.

Bud load is the most important factor affecting yield and cluster quality as well as vine vigor of Thompson seedless grapevines Morris and Cawthon (1980); Fawzi *et al.*, 1984; Marwad *et al.*, (1993); Omar and Abdel-kawi, 2000; Rubio *et al.*, (2002) on

Tempranillo variety Khamis *et al.*, (2017) on superior grapevines. Adjusting the length of fruiting spur is necessary to balance growth and fruiting status/vine nutritional status and bud fertility (Tamura *et al.*, 2002 and Ranspise *et al.*, 2003). Previous studies showed that adjusting the length of fruiting spurs in most grapevine cvs. is considered the main reasons for promoting productivity (Abdel-Fattah *et al.*, 1993; Kamel, 2002; Ansam, 2002; Awad, 2003; Nejatian, 2003; Jarad, 2004; Abdel-Mohsen, 2013; Ali and Moumen Kh. Mohamed, 2016) and Khamis *et al* (2017)

Therefore the objective of this study is to determine the optimum bud load for Early sweet grapevine and their effect on bud behavior, fruit quality, yield and vegetative growth.

### Materials and Methods

This study was carried out during two consecutive 2016 & 2017 experimental seasons in a private vineyard at El-Khatatba, Menoufiya governorate, Egypt on Early sweet grapevines cultivar. The vines were 7-year-old, grown in a sandy soil, spaced at 2 X 3 meters apart, irrigated by the drip irrigation system, trellised by the Spanish Parron shape system.

Vines were trained to quadrilateral cordon and spur-pruned. The experiment consisted of nine treatments arranged in a randomized complete block design, a hundred and eight uniform vines were chosen. Each twelve vines acted as a replicate and each three replicates acted as treatment. The vines were pruned during at the last week of December during two seasons of study. All vines received the same cultural managements recommended by ministry agriculture.

### The study included the following treatments

T1-pruning at 24 fruiting spurs ×2 buds= 48 buds  
 T2-pruning at 12 fruiting spurs ×4 buds= 48 buds  
 T3-pruning at 8 fruiting spurs ×6 buds= 48 buds  
 T4-pruning at 36 fruiting spurs ×2 buds=72 buds  
 T5-pruning at 18 fruiting spurs ×4 buds=72 buds  
 T6-pruning at 12 fruiting spurs ×6 buds=72 buds  
 T7-pruning at 48 fruiting spurs ×2 buds=96 buds  
 T8-pruning at 24 fruiting spurs ×4 buds=96 buds  
 T9-pruning at 16 fruiting spurs ×6 buds=96 buds

### 1-Bud behavior

During the two seasons, the following measurements were recorded:

#### a-Bud burst:

##### b- Percentage of bud burst:

Number of buds were counted one month after bud burst and the percentage of bud burst were calculated as follows according to **Bessis(1960)**.

$$\text{Bud burst\%} = \frac{\text{No of bursted buds per vine}}{\text{Total buds per vine}} \times 100$$

#### c-Bud fertility:

Number of clusters per vine were counted and divided by the total number of buds and the fertility was calculated as follows according to **Bessis(1960)**.

$$\text{Bud fertility\%} = \frac{\text{No of clusters per vine}}{\text{no of Total buds left at winter pruning}} \times 100$$

### 2-Morphological and vegetative growth

Vegetative growth parameters were determined after two weeks of fruit set as follows:

#### 1- Average shoots length (cm).

Twenty vegetative shoot were measured as average (cm), also number leaves fall date were calculated for each treatment when the leaves began take the red color and begins fall ( **El-Ashram 1993**).

#### 2- Average leaf area (cm<sup>2</sup>)

Twenty leaves / vine were picked at Veraison of the apical 6<sup>th</sup> and 7<sup>th</sup> leaves using a CI-203- Laser Area-meter made by CID, Inc., Vancouver, USA..

#### 3-N, P,K and Mgcontent in the leaves:

At full bloom, samples of 20 leaf petioles per each replicate were taken from leaves opposite to cluster were used for the determination of N, P and K according to (**Cotteniet al.1982**).

**4- Chlorophyll content in theleaves:**Sixth and seventh leaves from the tip of the growing shoots were used for the determination of total chlorophyll content in the leaves after two weeks from last treatment according to (**Mackinny, 1941**)

#### 3-Yield and physical characteristics of clusters

Harvesting indices (TSS% and acidity %) were weekly monitored from version till maturity when TSS reached about 16-17% according to **Tourk et al.,(1995)**.

- a- Average cluster length (cm).
- b- Average cluster width (cm).
- c- Average cluster weight (g).
- d- Yield /vine =number of clusters/vine × average cluster weight.

### 4-physical characteristics of berries

- a- Average berry weight (g).
- b- Total number of berries per cluster
- c- Average size of berries(cm)<sup>3</sup>
- d- Average berry width(cm)
- e- Average berry length(cm)
- f- Shot berries per cluster%=Number of shot berries per cluster ×100

Total number of berries per cluster

### 3- Chemical characteristics of berries

- a- Total soluble solids (TSS % in berry juice using a hand refract meter.
- b- Total treatable acidity (as tartaric acid %) according to the Official Analysis Methods (**A.O.A.C., 2000**).
- c- TSS / acid ratio.

### At dormant seasons:

- a- Wood ripening :
- b-Weight of pruning: was carried out at the time of winter pruning(g)
- c-Total carbohydrates in cane content (%):from fruitingcanes for next seasonwere determined calorimetrically by using reagent according to the method described by **Herbert et al.,( 1971)**

### Statistical analysis :

All the obtained data during ecal season of this study were subjected to statistical of various according to the method described by (**Snedecor and Cochran,1990**).However, the differences means were differentiated by using Duncan's multiple rage test (**Duncan,1955**).

### Result and Discussion

#### 1-Bud behavior

##### a-Time of bud burst:

Data presented in table (1) showed that pruning Early sweet grapevines atlevel 24spurs ×2 buds (48 buds/vine) advanced the beginning of bud burst date compared with other treatments followed by the vines were pruned at36 spurs ×2 buds (72buds/vine)and finally the vineswere pruning at level16 spurs ×6 buds (96 buds/vine) in both seasons of study.

##### b-Percentage of bud burst and bud fertility:

From table (1) it is clear that the bud burst and bud fertility percentage was increased significantly by decreasing the number of bud load/ vine in the two seasons. Since, the highest bud burst and bud fertility

percentage was associated with the lowest bud load (48 buds/vine) buds compared with (72 buds/vine) and (96 buds/vine) respectively in both seasons of study. Treatments under each levels (48, 72 or 96 buds) were nonsignificantly between them in both seasons. Moreover, vines were pruned at 8 spurs with 6 buds gave the highest bud burst and bud fertility percentage followed by the vines were pruned at 12 spurs  $\times$  4 buds compared with the other treatments while vines were pruned at 24 spurs with 4 buds gave the lowest bud

burst and bud fertility percentage compared with the other treatments during both seasons under the study.

The results obtained also by **Ali and Moumen Mohamed, 2016, Omar and Abd EL-Kawi (2000)** found that increasing bud load reduces bud fertility%. Furthermore, leaving 18 canes with 4 nodes and leaving 21 canes with 3 nodes give a highest significant value of bud fertility percentage in the first and second seasons respectively while the lowest value was obtained when leaving 8 cane with 6 nodes in both season of the study.

**Table 1.** Effect of different buds load levels on bud behavior of Early Sweet grapevines in 2016 and 2017 seasons

Treatments	Bud burst date		Bud burst%		Bud fertility%		
	2016	2017	2016	2017	2016	2017	
48	24spurs $\times$ 2 buds	13/2	12/2	96.2a	97.5a	78.3a	81.2 b
	12 spurs $\times$ 4 buds	15/2	14/2	96.9a	98.45a	82.8a	89.6a
	8 spurs $\times$ 6 buds	15/2	14/2	96.9a	98.5a	82.3a	91.2a
72	36spurs $\times$ 2 buds	14/2	15/2	91.3ab	93.2ab	57.4b	57.9c
	18 spurs $\times$ 4 buds	16/2	17/2	90.1ab	91.8ab	56.9b	57.3c
	12 spurs $\times$ 6 buds	17/2	15/2	89.9ab	91.7ab	54.2b	58.0c
96	48 spurs $\times$ 2 buds	17/2	16/2	87.5ab	90.0b	40.9c	41.9 d
	24 spurs $\times$ 4 buds	18/2	18/2	84.0b	87.45b	38.0c	47.2d
	16 spurs $\times$ 6 buds	20/2	19/2	87.5ab	89.6b	47.9b	41.9d

Values within each column followed by the same letter/s are not significantly different at 5% level

## 2- Vegetative growth

### a- Shoot length and Leaf area

With respect to the effect of buds load on shoot length and leaf area data obtained in Table (2) show that 2 buds/spur treatments was the superior treatment in this respect as it enhanced shoot length and leaf area in Early sweet grapevines as compared with 4 and 6 buds/spur treatments in this concern during 2016 and 2017 seasons.

The vine were pruning at 24 spurs/2 buds (48 buds/vine) and 48 spurs/2 buds (96 buds/vine) gave the highest value compared with the other treatments during both seasons under the study, However The vine were pruning at 12 spurs/8 buds (96 buds/vine) recorded the lowest shoot length and leaf area in both seasons.

This increased with sever pruning may be due to the strong reduction in pruned wood as compared to light pruning, thus affects that balance between nitrogen and carbohydrates in favorite of nitrogen which promotes vegetative growth. The findings of **Abd El-Baki (2003)** showed that there was a gradual increase in shoot length of both "King Ruby" and "Thompson seedless" all along the growing seasons. However, "King Ruby" vines produced the longest shoots under the level of 12 buds/vine followed by 18 and 24 buds/vine, while the shortest ones were noticed in case of 30 buds/vine. In addition, **Ali and MoumenKh. Mohamed, (2016)** found that the highest shoot length and leaf area were recorded on the vines that pruned to leave 15 fruiting spurs  $\times$  4eyes. Also,

**Sabbatiniet al. (2015)** mentioned that as number of nodes retained increased, vine size and leaf area.

### b- chlorophyll A and B

Regarding chlorophyll A and B data in Table (2) revealed that pruning Early sweet grapevines at level 24spurs  $\times$  2 buds (48 buds/vine) increased chlorophyll A and B followed by the vines were pruned at 12 spurs  $\times$  4 buds (48 buds/vine) as compared with the other treatments in 2016 and 2017 seasons, respectively. While, vines pruned at 48 spurs with 2 buds (96 buds/vine) gave the lowest chlorophyll A and B compared with the other treatments in both seasons of study.

These results are in contrast with those reported by **Velu (2001)** and **Senthilkumar et al (2015)** observed that the maximum chlorophyll content (2.699 mg/g) was registered at a pruning level of 67 % of canes to 5 bud level and 33 % of the canes to 2 bud level. In addition, **Ali, et al., (2016)** found that the highest chlorophyll A and B were recorded on the vines that pruned to leave 8 canes  $\times$  10 eyes

**Table 2.** Effect of different of buds load levels on Shoot length, Leaf area chlorophyll A and B of Early Sweet grapevines in 2016 and 2017 seasons

Treatments	Shoot length (cm).		Leaf area (cm <sup>2</sup> )		chlorophyll A (mg / g f.w.)		Chlorophyll B (mg / g f.w.)		
	2016	2017	2016	2017	2016	2017	2016	2017	
48	24 spurs ×2 buds	136.0a	139.0a	192.2a	200.9a	1.62a	1.69a	1.52a	1.51a
	12 spurs ×4 buds	131.3b	134.3bc	178.5ab	182.8ab	1.61a	1.75a	1.51a	1.16a
	8 spurs ×6 buds	129.5b	132.8c	92.8c	101.1c	0.94bc	1.93a	0.86	1.16a
72	36spurs ×2 buds	132.6b	136.8abc	191.9a	195.9a	0.95bc	0.96bc	0.85bc	0.45b
	18 spurs ×4 buds	130.3b	133.5c	175.5ab	183.8ab	0.82bc	0.96bc	0.95bc	0.44b
	12 spurs ×6 buds	129.8b	133.3c	88.1c	97.3c	0.94bc	0.95bc	0.93bc	0.46b
	48 spurs ×2 buds	135.8a	137.8ab	192.1 a	199.8a	0.55c	0.57c	0.50 c	0.38b
96	24 spurs ×4 buds	131.0b	134.8bc	177.1ab	180.9ab	1.56a	0.75bc	0.72bc	0.41b
	16 spurs ×6 buds	129.5b	133.8bc	97.0c	103.8c	0.80bc	0.80bc	0.76	0.33b

Values within each column followed by the same letter/s are not significantly different at 5% level

### N, P, K and Mg content in the leaves

The data presented in Table 3 revealed that vines were pruned at 24 spurs/2 buds (48 buds/vine) gave the highest percentage of phosphorous, potassium and magnesium compared with (72 buds/vine) and (96 buds/vine) respectively in both seasons of study followed by the vines were pruned 12 spurs ×4 buds as compared with the other treatments in 2016 and 2017 seasons, respectively. On the other hand, the lowest value percentage of phosphorous, potassium and magnesium was obtained when the vine were pruning at 48 spurs ×2 buds (96 buds/vine) in both seasons of the study while different levels of buds load gave a non significantly differences in percentage of nitrogen compared of them under this study.

These results as a general trend are in agreement with (Weaver, 1976) The increase on the leaf chemical constituents at the higher levels of vine load might be attributed to the promotion on the leaf area which resulted in enhancing photosynthesis process as well as the reduction on main shoot length and number of leaves which aids in reducing the depletions of these nutrients. Ali, (2016) showed that the maximum values of chlorophylls (a & b), total chlorophylls, N, P, K and Mg in the leaves were recorded on the vines pruned to leave 102 eyes/vine and sprayed three times with citric acid at 0.4%. These findings were true during both seasons.

**Table 3.** Effect of different levels of buds load on N, P, K and Mg of Early Sweet grapevines in 2016 and 2017 seasons

Treatments	N %		P %		K %		Mg %		
	2016	2017	2016	2017	2016	2017	2016	2017	
48	24 spurs ×2 buds	1.88a	1.58a	0.36a	0.39a	0.86a	0.89a	0.49a	0.50a
	12 spurs ×4 buds	1.61a	1.53a	0.36a	0.38a	0.95a	0.95a	0.49a	0.49 a
	8 spurs ×6 buds	1.43a	1.73a	0.36a	0.37a	0.84a	0.85a	0.48a	0.48a
72	36spurs ×2 buds	1.46a	1.54a	0.31a	0.34a	0.64bc	0.66bc	0.36b	0.36b
	18 spurs ×4 buds	1.56a	1.26a	0.29a	0.35a	0.69b	0.70b	0.34b	0.35 b
	12 spurs ×6 buds	1.39a	1.75a	0.30a	0.32a	0.57cd	0.58cd	0.33b	0.33b
	48 spurs ×2 buds	1.69a	1.36a	0.18b	0.20b	0.51d	0.53d	0.31b	0.31b
96	24 spurs ×4 buds	1.64a	1.61a	0.19b	0.21b	0.51d	0.53d	0.31b	0.32 b
	18 spurs ×6 buds	1.45a	1.57a	0.19b	0.21b	0.50d	0.31 b	0.31b	0.32b

Values within each column followed by the same letter/s are not significantly different at 5% level

### 3-Yield and physical characteristics of clusters

#### a- Cluster length and width

Data in Table (4) declared that the buds load of (48 buds/vine) 24 spurs/2 buds produced the highest significant cluster length and width compared with pruning (72 buds/vine) and (96 buds/vine) in both seasons. However the treatment of (96 buds/vine) 24 spurs/4 buds recorded the lowest cluster length and width in both 2016 and 2017 seasons

#### b- Cluster weight and Yield /vine

data in Table (5) revealed that pruning Early sweet grapevines at (48 buds/vine) 24 spurs/2 buds produced the highest cluster weight compared with pruning at (72 buds/vine) and (96 buds/vine) in both seasons. While, pruning at (72 buds/vine) 12 spurs/6 buds recorded the lowest cluster weight in both 2016 and 2017 seasons. Regarding yield per vine pruned (96 buds/vine) 24 spurs/4 buds produced the highest yield per vine compared with leaving (72 buds/vine) and (48 buds/vine) in both seasons. On the other hand, pruning

at (48buds)24 spurs/2 buds recorded the lowest yield per vine in both 2016 and 2017 seasons.

These results agreed with those findings of **Ansam(2002)and EL-Bazet al., (2002)** On Kings Ruby grapevine vines pruned to 40 or 60 buds/vine produced the highest width and length of clusters

compared with the other bud load (20, 50 and 80bud/vine) **Aly (2001).Fawziet al. (2015)** showed that interaction between the two studied factors was insignificant in most cases. The highest value of cluster width and cluster length was noticed with the pruned vine at 2eyes per cane plus 24buds.

**Table 4.** Effect of different of buds loadlevels on Cluster length (cm), width (cm ), Cluster weight (g) and Yield /vine (kg) of Early Sweet grapevines in 2016 and 2017 seasons

Treatments	Cluster length (cm)		Cluster width (cm)		Cluster weight (g)		Yield /vine (kg)	
	2016	2017	2016	2017	2016	2017	2016	2017
24 spurs ×2 buds	28.3a	29 a	24.2a	25.8a	512.5 a	525 a	15.75 d	17.55 c
48								
12 spurs ×4 buds	27.5a	29 a	24.8a	23.8 ab	412.5bcd	445 bc	17.01bcd	18.91abc
8 spurs ×6 buds	27.4a	28.a	16.5 de	15.3 e	385 d	410 cd	17.35bcd	19.03abc
36spurs ×2 buds	24 b	26ab	22ab	22.3abc	400 cd	437.5 bc	18 bc	20.23ab
72								
18 spurs ×4 buds	23.5b	24.5b	19.5bcd	20.3bd	387.5 d	387.5cd	19.01 ab	19.5 abc
12 spurs ×6 buds	27.8a	27.3b	19.8bcd	18.8cde	375 d	367.5 d	16.35 cd	18.45 bc
48 spurs ×2 buds	23.8b	24.5b	20.5bc	19.8bce	460 abc	497.5ab	18.37abc	20.56 a
96								
24 spurs ×4 buds	22 b	23 b	15.75 e	16.8de	462.5ab	512.5 a	20.21 a	20.67 a
16 spurs ×6 buds	22.8b	24.3b	18cde	19 cde	487.5 a	522.5 a	18.23 abc	19.93 ab

Values within each column followed by the same letter/s are not significantly different at 5% level

#### Physical characteristics of berries

Data presented in table (5) revealed that the highest values of berry weight, size, width and length were obtained when the vines were pruned at(48 buds/vine) 24 spurs/2 buds as compared with the other treatments in 2016 and 2017seasons followed by the vines were pruned at (48 buds/vine) 8 spur ×6 buds as compared with the other treatments in 2016 and 2017seasons, respectively. While, vines were pruned at(96 buds/vine) 24 spurs with 4 buds gave the lowest berry

weight, size, width and length compared with the other treatments in both seasons of study.

Regarding shoot berries percentage per cluster of in the same table revealed that pruning Early sweet grapevines (48 buds/vine) 12 spurs/4 buds decreased shoot berriespercentage per cluster as compared with(72 buds/vine)and (96 buds/vine) in 2016 and 2017seasons, respectively. While, vines pruned to (96 buds/vine) 48 spurs with 2 buds gave the highest shoot berriespercentage per cluster of compared with the other treatments in both seasons of study.

**Table 5.** Effect of different of buds load levels on 25 berries weight (g), Size of berries (cm)<sup>3</sup>, Berry width, length(cm)and shoot berries per cluster%, of Early Sweet grapevines in 2016 and 2017 seasons

Treatments	25 berries weight(g)		Size of berries(cm) <sup>3</sup>		Berry width(mm)		Berry length(mm)		shoot berries per cluster%	
	2016	2017	2016	2017	2016	2017	2016	2017	2016	2017
24 spurs ×2 buds	117.2a	120.a	115.4a	118.a	19.8a	21a	23.2a	23.7a	2.8e	3.1d
48										
12 spurs ×4 buds	96bc	99.3 bc	102.5 bcd	108.3 b	19 ab	19.6 ab	22.6 a	23.1 a	2.1 e	2.6 D
8 spurs ×6 buds	101.3 b	105 b	105 bc	108 b	19.5 ab	20.3 ab	22.4 a	22.9 a	3.7 cde	4.1 Cd
36spurs ×2 buds	86.7 Bcd	93.3 bcd	97.7 bcd	100.7 b	19.7 ab	18.9 abc	23 a	23.7 a	6.2 bcde	7.4 Bc
72										
18 spurs ×4 buds	96.8 Bc	103 b	99.5 bcd	102.3 b	17.7 bc	18.4 bc	19 bc	20 bc	9.4 b	10.1 B
12 spurs ×6 buds	82.8 Cde	87.5 cde	96.3 cd	100 b	18.7 ab	19.2 abc	21.6 a	22.1 ab	6.9 bc	7.5 Bc
48 spurs ×2 buds	73.5d ef	81 de	95 d	100.3 b	18.4 ab	18.9 abc	21.4 a	22 ab	19.1 a	19.7 A
96										
24 spurs ×4 buds	66.75 f	77.5 e	61.3 f	77.5 c	16.5 cd	17.4 cd	17.6 c	18.4 c	17.2 a	18.0 A
16spurs ×6 buds	70 ef	75 e	75 e	80 c	15.9 d	16.2 d	18.1 c	18.7 c	16.4 a	17.4 A

Values within each column followed by the same letter/s are not significantly different at 5% level

Ali and MoumenKh. Mohamed, 2016 showed that the highest berry weight, size, width and length were recorded on the vines that pruned to leave 15 fruiting spurs x 4eyes and decreasing shoot berries per cluster percentage of Early Sweet grapevines. Also, Hussain and El-Dujaili, (1990), Abd El-Baki (2003) and Sabbatiniet al. (2015) mentioned that as number of nodes retained increased, berry weight of 'Niagara' grapevines increased

### Chemical characteristics

#### Total soluble solids content (%), Acidity and TSS/acid ratio

Data presented in Table (6) indicated that non significantly difference between all treatments in both seasons on total soluble solids content (%), Acidity and TSS/acid ratio of Early sweet

grapevinescultivar.Since,vines were pruned at (48 buds/vine) 12 spur x4 buds and (48 buds/vine) 24spur x2 buds gave the highest total soluble solids content and decreased Aciditycompared with treatments (72 buds/vine)and (96 buds/vine) respectively in both seasons of study while treatment (96 buds/vine) 48 spurs x2 buds gave the lowest value oftotal soluble solids contentand increased Aciditycompared with the other treatments in 2016 and 2017seasons this study

This is agreement with those mentioned by Dhillon (2004), Almanza-Merchan et al. (2014) and Sabbatini et al. (2015) stated that long pruning type presented the highest values of TSS%. Beside, Rizket al. (1994) and Abd El-Wahab (1997) recorded that total soluble solids of grape berries was not affected by treatments of bearing unit length. In this trend, Ansam (2002) and Cangi and Klc (2011) found that bud loading had no effect on total soluble solids in addition.

**Table 6.** Effect of different of buds loadlevels on TSS, Acidity% and TSS/acid ratio of Early Sweet grapevines in 2016 and 2017 seasons

Treatments	TSS%		Acidity%		TSS/acid ratio		
	2016	2017	2016	2017	2016	2017	
48	24 spurs x2 buds	17.2 a	17.7a	0.374 a	0.374ab	45.9a	45.7 a
	12 spurs x4 buds	17.3 a	17.7a	0.363b	0.375 a	47.6a	47.2 a
	8 spurs x6 buds	16.8 ab	17.3ab	0.367ab	0.375 a	45.4ab	46.1 a
72	36spurs x2 buds	16ab	16.5ab	0.373ab	0.387a	42.8ab	42.6ab
	18 spurs x4 buds	16.4 ab	16.8ab	0.365ab	0.375a	44.9ab	44.8ab
	12 spurs x6 buds	16.3 ab	16.8ab	0.367 ab	0.378a	44.4ab	44.4ab
96	48 spurs x2 buds	16.5a b	16.0ab	0.387a	0.390 a	42.6.1ab	41.0b
	24 spurs x4 buds	16.8 ab	16.1ab	0.373ab	0.383a	45.0ab	42.0ab
	16 spurs x6 buds	16.8 ab	16.0ab	0.375a	0.365ab	44.0ab	43.8ab

Values within each column followed by the same letter/s are not significantly different at 5% level

### At dormant seasons

#### Carbohydrates, Wood ripening and Weight of pruning/vine

The data presented in Table (7) revealed vines were prunedat(48 buds/vine)24 spurs/2 buds gave the highest Wood ripening and Weight of pruning per vinecompared with (72 buds/vine)and (96 buds/vine) respectively in both seasons of study.On the other hand ,the lowest value ofwood ripening and weight of pruning per vine was obtained whenvines were pruning at (72 buds/vine)12spurs x6 buds in both seasons of the study while different levels of buds load gave a non significantly differences in carbohydrates in canes except pruning (72 buds/vine)12 spur x6 buds gave the lowest valuecompared with other treatments of study. Our results in this connection agree with those obtained by Omar & Abdel-kawi (2000) who

reported that increasing leaves lead to heavy canopy with increase in active photosynthesis and stored carbohydrates in the new canes. Similar results were obtained by El-Bazet al., (2002) on Crimson seedless grapevines Genaidy (2015) increasing the buds on the vineyard led to a weight loss of wood pruning and mature wood.

### Conclusion

From the previous results, it can be recommended that pruning Early Sweet grapevine at (48 bud/vine) namely 24,12 and 8 fruiting spurs adopted to 2,4and 6 buds/spur gave highest values of bud behavior, physical characteristics and vegetative growth .Also increased wood ripening ,weight of pruning and total carbohydrate in canes However, the bud load of (96 bud/vine) 24spurs x4 buds gave the highest with yield per vine in both season in both seasons

**Table 7.** Effect of different of buds load levels on carbohydrates, Wood ripening and Weight of pruning/vine(g) of Early sweet grapevines in 2016 and 2017 seasons

Treatments	carbohydrates(%)		Wood ripening		Weight of pruning/vine(g)		
	2016	2017	2016	2017	2016	2017	
48	24 spurs ×2 buds	35.56 a	35.60 a	0.962a	0.969a	3.77a	4.13a
	12 spurs ×4 buds	31.10 ab	35.07 a	0.903b	0.912b	2.83b	3.07b
	8 spurs ×6 buds	31.94 ab	34.71 a	0.813c	0.826c	2.70b	2.91b
72	36spurs ×2 buds	34.91 a	31.16 ab	0.957a	0.966a	3.96a	4.04a
	18 spurs ×4 buds	32.70 a	27.86 b	0.902b	0.910b	2.90b	3.125b
	12 spurs ×6 buds	31.94 ab	32.02 ab	0.755d	0.770d	2.60b	2.85b
96	48 spurs ×2 buds	34.96 a	34.98 a	0.947ab	0.936ab	3.75a	3.94a
	24 spurs ×4 buds	34.65 a	32.745 a	0.903c	0.911b	3.06b	3.28b
	16 spurs ×6 buds	27.7 b	33.88 a	0.770d	0.779d	2.73b	3.025b

Values within each column followed by the same letter/s are not significantly different at 5% level

## References

- A.O.A.C.(2000). Association of Official Agricultural Chemists. Official Methods of Analysis. 4th ed.pp. 495-510. Benjamin Franklin Station, Washington. D.C., U.S.A.
- Abd El-Baki, M.S.A. ( 2003). Effect of summer pruning and thinning on growth and fruiting of King Ruby grapes. M.Sc. Thesis., Fac. of Agric., Cairo Univ., Cairo, Egypt
- Abd El-Wahab, M. A. (1997).Effect of cane length on bud behaviour, growth and production of "king Ruby" grapevines.M. Sc. Thesis, Cairo University, Egypt.
- Abdel-Fattah S.E; Marwad, I. A; and Isis AbdelshahiedRizk (1993). Effect of bud load and spur length on Roumi Red grapevine I-Weight of pruning and chemical composition of 1 year - old wood. Zagazig.J. Agric.. Res Vol 2.0 No. (6) pp1889 -1899
- Abdel-Mohsen, M.A. (2013): Application of various pruning treatments for improving productivity and fruit quality of crimson seedless grapevine. World Journal of Agricultural Sciences 9 (5): 377-382
- Almanza-Merchan, P.J., G.R. Fischer and G.E. Cely( 2014).The importance of pruning to the quality of wine grape fruits (*Vitisvinifera* L.) cultivated under high-altitude tropical conditions. *Agronomia Colombiana*, 32(3): 341-348.
- Aly, M.S.M.( 2001) Effect of bud load and some growth regulators on growth, yield and fruit quality of King's Ruby grapevine cultivar. Ph. D. Agric. Sci., Ain Shams University, Cairo, Egypt
- Ali, A.H.; M.Kh. Uwakiem and H.M.M. Sayed(2016)Effect of Vine Load and Spraying Citric Acid on Fruiting of Superior Grapevines Grown Under Minia Region Conditions- Egypt *Assiut J. Agric. Sci.*, (47) No. (6-2) 2016 (484-503)
- Ali H. Ali and Moumen A. Kh. Mohamed( 2016) Effect of Fruiting Spur Length and Spraying Seaweed Extract on Yield and Berries Quality of Early Sweet Grapevines *Assiut J. Agric. Sci.*, (47) No. (6-2) 2016 (504-517) ISSN: 1110-0486
- Ansam, S.A. El.,(2002). Effect of cane length on bud behaviour, growth and productivity of "Superior" grapevines. M.Sc. Thesis, Fac. of Agric., Cairo Univ., Egypt.
- Awad, N.A.E. (2003): Studies on pruning severity of Thompson seedless grapes. M. Sc. Thesis Fac. of Agric. Mansoura Univ. Egypt
- Bessis, R. (1960). Sur Differentes Models Expression Quantitative Delafertilité chez la vigneAca p.p. 828 -882.
- Cangi, R. and D. Klc, (2011) Effects of bud loading levels and nitrogen doses on yield, physical and chemical properties of brined grape-leaves. *African Journal of Biotechnology*, 10(57): 12195-12201.
- Celik, H., Y.S. Agaoglu, Y. Fidan, B. Marasali and C. Soylemezoglu, (1998). General Viticulture. Professional Books Series, Ankara, 1: 1-253.
- Cottenie, A.; M. Verloo; L. Kiekens; G. Relgho and W. Camerlynck (1982). Chemical analysis of plant and soil. Lab. of analytical and Agrochemistry State Univ. Gent, Belgium
- Dhillon, W.S. (2004).Standardization of pruning intensity in grape cv. Perlette. *Haryana Journal of Horticultural Sciences*, 33(3/4): 172-174.Cangi, R. and D. Klc, 2011. Effects of bud loading levels and nitrogen doses on yield, physical and chemical properties of brined grape-leaves. *African Journal of Biotechnology*, 10(57): 12195 -12201.
- Duncan, B. D. (1955). Multiple test range and multiple F tests. *Biometrics*. 11-142.
- El- Ashram, M.A. (1993): Modern Approaches of Cultivation and Production of Grapevines. Dar El-Fekr, El- Arabie, pp. 275-276.
- EL-Baz, El. El. T.; A.M. Mansour; El.F.El-Dengawy and B.N. Samra (2002) Influence of pruning severity on bud behavior, yield, ,berry quality and some biochemical contents of the canes of "Crimson seedless" Grapes. *Egypt.j.Hort.*29(1) pp.39-60.
- Fawzi, F., Bondok, A.Z. and Ghobrial, G. F. (1984). Effect of cane length on bud behaviour and wood ripening of Thompson seedless grape variety,

- Annals Agric., Sci., Ain Shams Univ., Cairo 29 (1) 465-474.
- Fawzi, F., F.H. Laila, M.F.M. Shahin, M.A. Merwad and E.A.E. Genaidy (2015). Effect of vine bud load on bud behavior, yield, fruit quality and wood ripening of superior grape cultivar. International Journal of Agricultural Technology, 11(5): 1275-1284.
- Genaidy, E. (2015). Effect of vine bud load on bud behavior, yield, fruit quality and wood ripening of superior grape cultivar. J of Agr. Tech, 1275-1284
- Herbert, D.; P.J. Phipps and R.F. Strange (1971). Determination of total carbohydrates Methods in Microbian; 5(B): 209-244.
- Hussain, M.M. and J.A.H. El-Dujaili (1990). Effect of bud level and training system on yield and quality of Dies El-Anz grape cultivar (*Vitis vinifera* L.). Hort. Abst., 62: 6472.
- Jarad, A. (2004): Effect of fruit canes and eyes vine load on the productivity of a local grape variety under the conditions of Deir El- Zor city. Bull Fac. Agric. Cairo Univ, 55: 603-608
- Kamel, M.K. (2002): Physiological studies on pruning and fertilization of Flame seedless grapevines. (*V. vinifera* L.). Ph. D. Thesis. Fac. Of Agric. Minia Univ. Egypt
- Khamis, M. A.; A.A.R. Atawia; H.E.M. El-Badawy and A.A.M. Abd EL-Samea (2017) Effect of buds load on growth, yield and fruit quality of Superior Grapevines. Middle East. J. of Agric. Res ISSN 2077-4605. Vol: 061 Issue: 01. J- Mar pag: 152-160
- Mackinny, G. (1941). Absorption of light by chlorophyll solution. J. Bio. Chem., 140: 315-322.
- Marwad, I. A.; Rizk, N. A and Ibrahim, A. H (1993). Effect of cane length on bud behaviour, Yield, bunch characteristics and wood maturity of Thompson seedless grapevines Egypt. J. Appl. Sci., 8 (12) pp 47-60.
- Morris, J. R. and D. L. Cawthan (1980). Yield and quality response of "Concord grapes to training systems and pruning severity in Arkansas, J. Amer. Soc. Hort. Sci., 105 (3): 307-10. (Hort. Abst. 51: 233).
- Nejatian, M.A. (2003): Effects of bud number and cane length on bud fertility and some characteristics of grapevine cv. Bidanehsefid, 19(4): 457-467.
- Omar, A.H and Abdel- kawi, A. (2000). Optimal bud load for Thompson seedless grapevines. J. Agric Sci. Mansoura Univ. 25 (9), pp 5769-5777.
- Ranspise, S. A.; Patil, B. T.; More, T. A.; Birade, R. M.; Ghure, T. K., (2003): Effect of sub-cane pruning on fruitfulness and yield of grape cv Thompson seedless. Journal of Maharashtra Agricultural Universities. 27 3: 258-259
- Rizk, I. A. (1996). Effect of cane length on bud behaviour and bunch characteristics in "Thompson seedless" cultivar. Egypt. J. Appl. Sci., 11 (7): 220-34.
- Rizk, N. A.; I. A. Rizk and V. H. Girgis (1994). Effect of cane length on bud behaviour, wood ripening and bunch characteristics in "Thompson seedless" grapevines. Egypt. J. Appl. Sci., 9 (5): 74-89.
- Rubio, J.A., Albuquerque M. V., peixy, J.L and yuste . J. (2002). la mejora varietal de la vid, sus consecuencias y beneficios. Vida Rural, pp 149: pp 62-66
- Sabbatini, P., K. Wierba, L. Clearwater and G.S. Howell (2015). Impact of training system and pruning severity on yield, fruit composition, and vegetative growth of 'Niagara' grapevines in Michigan International Journal of Fruit Science, 15(3): 237-250.
- Senthilkumar, S, Vijayakumar, R.M., Soorianathasundaram, K. and Durgadevi, D. (2015) Effect of pruning severity on vegetative, physiological, yield and quality attributes in grape (*Vitis vinifera* L.) : A Review. Curr. Agri Res. J. Vol. 3(1), 42-54.
- Snedecor, G.A. and W.G. Cochran (1990). Statistical Methods. 6<sup>th</sup> ed the Iowa state. Univ. press. Iowa U.S.A.
- Tamura, F.; Muraya, K. and Fujji, Y. (2002): Effects of short cane pruning in summer and long cane pruning in winter on growth and yield of the grapevine "Muscat" of Alexandria' in forcing culture condition. Hort. Res. (Japan), 1(4): 269-274. (Hort. Abst., 73(5): 4269).
- Tourky, M.N., El-Shahat, S.S. and Rizk, M. H. (1995). Effect of Dormex on fruit set, quality and storage life of Thompson seedless grapes (Banati grapes) J. Agric. Sci., Mansoura Univ., 20(12): 5139-5151.
- Velu, V. (2001) Studies on bud load and certain crop thinning practices on vigour, yield and quality of grapes (*Vitis vinifera* L.) cv. Muscat. M.Sc.,
- Weaver, R.J. (1976). Grape Growing. Wiley-Interscience Publication, John Wiley & Davis, New York, London, Sydney, Toronto pp. 160-175.

## تأثير حمولة البراعم على سلوك البراعم وجودة الثمار في العنب الايرلى سويت

ثرىا صابر على ابو الوفا

قسم بحوث العنب -معهد بحوث البساتين -مركز البحوث الزراعيه

أجريت هذه الدراسة خلال موسمي ٢٠١٦ و ٢٠١٧ في مزرعة خاصة بالخطاطبه- محافظة المنوفيه كرمات عنبايرلى سويت عمرها ٧ سنوات ومنزعة في تربة رملية تروى بالتنقيط وعلى مسافة زراعة ٣×٢ م ومرياه بطريقة الكردون الرباعي باستخدام نظام تدعيم البارون سيستم . وقد استهدف هذا البحث دراسة تأثير حمولة البراعم على سلوك العيون وجودة الثمار في العنب الايرلى سويت. حيث تم استخدام ثلاث مستويات مختلفة من حمولة البراعم وهي (٤٨-٧٢-٩٦ برعم/للكرمه) عدد البراعم لكل دابره ثمرية (٢-٤-٦) وقد أظهرت النتائج أن تحميل ٤٨ برعم لكل كرمه اى ٢٤ دابره ثمرية على كل دابره ٢ عين كان له تأثير كبيرسلوك البراعم و الصفات الطبيعية و الكيميائية للعنقود والحباب وتحسن صفاتالنمو الخضري والهرمونات فى الأوراق وتحسين نسبة الكربوهيدرات فى القصباب . ومع ذلك كانت أفضل المعاملات إنتاجا للمحصول ٩٦ برعم لكل كرمه وفى النهاية كانت المعاملة ٤٨ برعم لكل كرمه اى ٢٤ و١٢ و٨ دوابر ثمرية مع ٢ و٤ و٦ براعم على الدابره الثمرية على التوالي تحت الدراسة افضل المعاملات