

Chemical Evaluation of Some Stevia (*Stevia rebaudiana*, Bertoni) Varieties of under Different Nitrogen Fertilization Levels

. A .A. Tantawy * .A.E. Attia **Nareyman. N. Kenawy. **.

*Agronomy Dept., Fac. of a Agric., Minia Universtity. **Agricultural Research Center, Giza, Egypt.

Correspondence author:nareyman33@gmail.com

Abstract

Two field experiments were carried out at agricultural research center, Giza, Egypt during 2015/2016 and 2016/2017 seasons to study the effect of nitrogen fertilizer levels (20, 30 and 40 kg N/ fed/cut.) on some stevia varieties (china1, Egy1, spanti and shou2). The highest nitrogen rate of 40 kg. N /fed /cut was significant in all studied yield and chemical characters and gave the highest values of stevioside%. recorded (6.09 and 6.62%), rebaudioside A% (7.23 and 8.04%), Stevioside kg /fed (238.9 and 286.1 kg) and rebaudioside A kg /fed (287.2 and 350.5 kg), respectively as compared with the other two nitrogen treatments in both seasons.

Stevia Varieties were significantly different in all studied yield characters. china 1 Varieties gave the highest values of stevioside% (6.83 and 7.34 %), rebaudioside A% (8.44 and 9.26 %), rebaudioside A kg /fed(300.8 and 363.2 kg), respectively of all studied yield and chemical characters except for dry leaves yield kg/fed where Egy1 varieties gave the highest values in the first and second seasons.

.Concerning the effect of the interaction between nitrogen fertilizer levels and stevia varieties on yield and chemical characters, nitrogen fertilizer level of 40 kg N/ fed/cut applied to china1 varieties gave the highest values in all studied yield and chemical composition except for dry leaves yield (kg/fed) where Egy1 varieties and nitrogen fertilizer level of 40 kg N/ fed/cut gave the highest values in the first and second seasons.

Key words: stevia, nitrogen fertilizer, **varieties**, stevioside, rebaudioside -A .

Introduction

Stevia is a new crop in Egypt. It was approved in 1995 for use as a food ingredient by Food Standard for Egypt. This plant is believed to be the most ideal substitute for sugar. Steviol Glycoside (SG) found in the leaves is responsible for the sweetness of stevia. As a new crop to for Egypt, commercial cultivation of stevia is yet to commence.

. Leaves of *Stevia rebaudiana* contains diterpene glycosides which taste sweet but with zero calories and 100–300 times sweeter than sugar (sucrose) (Megeji *et al* 2005 and Soejarto *et al* 1983).

This plant is believed to be the most ideal substitute for sugar and important to assist in medicinal value. Given its many health benefits, recent GRAS acceptance and existing evidence in support of stevia's safety for human consumption. namely stevioside, rebaudioside A, B, C, D and E (Geuns 2003). The plant is known to exhibit a wide range of biological activities like hypoglycemic, anti-oxidant, anticancer, antibacterial activities. Stevia contains a high percentage of phenols, flavonoids and antioxidant activity (Negar *et al* 2012) . *Stevia rebaudiana*, Bertoni leaves are a natural source of diterpenic glycosides, and various bioactive compounds. characterize antioxidants (Danijela *et al* 2018). Stevia now is used as sweeteners to replace sugar in foods, beverages and medicines, stable at wide range of temperature and pH conditions during processing and offer no shelf life limitations,(Balwinder *et al* 2014). They are thermostable even at temperatures of up to 200 !C,

making them suitable for use in cooked foods.(Roberto *et al* 2012).

Materials and methods

The present work was designed to study the respons of (yield and chemical properties of some stevia (*Stevia rebaudiana*, Bertani) varieties under different nitrogen fertilization levels. Two experiments were carried out at Agricultural Research Center , Giza, Egypt during 2015/2016 and 2016/2017 seasons.

Treatments: Two factors were studied as follow:

Factor A: nitrogen fertilizer levels (N) : N1= 20 kg N fed/cut, N2:=30 kg N fed/cut and N3= 40 kg N fed/cut.

Factor B: Varieties (v): V1: china 1, V2 : Egy 1, V3 : spanti, V4 :shou 2.

Nitrogen fertilizer was applied in the form of urea (46.5% N) in two equal rates, the first rates was applied after 15 days of planting (or after cutting in the following cuts) and second rates was applied after 15 days from the first rates.

Field Experimental Design:

The experimental design used was a split plot design with three replicates, Fertilizer levels were arranged in the main plots and cultivars in the sub plots. The sub plot area was 10.8 m² (6 ridges) 3.0 meters in the length and 60 cm in width The treatments (12) were the combinations of three nitrogen levels and four varieties

Table (A) Physical and Chemical analysis of experimental soils in 2015/2016 and 2016/2017 seasons.

Analysis	Seasons	
	2015/2016	2016/2017
Mechanical analysis		
Coarse sand %	0.20	0.50
Fine sand %	12.16	14.2
Silt %	48.85	45.7
Clay %	38.99	36.5
Textural class	Cilt clay loam	Cilt clay loam
Chemical analysis		
Organic matter %	1.08	1.32
CaCo ₃ %	1.10	2.0
PH(1:2.5)**	8.10	8.0
Soluble Ca ⁺⁺ (meq /100 g soil)	11.2	16.2
Soluble Mg ⁺⁺ (meq /100 g soil)	10.7	14.7
Soluble Na + (meq /100 g soil)	8.3	11.7
Soluble K + (meq /100 g soil)	4.1	2.6
Soluble Co ₃ --(meq /100 g soil)	--	--
Soluble Hco ₃ --(meq /100 g soil)	3.5	5.6
Soluble Cl--(meq /100 g soil)	13.3	17.9
Soluble So ₄ --(meq /100 g soil)	17.3	31.7
Available N(ppm)	11.00%	13.9%
Available P(ppm)	9.12%	11%
Available K(ppm)	35.86%	34.1%

Studied characters

- Dry leaves yield kg/fed/ year
- Stevioside%-
- Rebaudioside A% -
- Stevioside yield (kg / fed / year) -
- Rebaudioside A yield (kg / fed / year)

Statistical analysis:

All collected data were statistically analyzed according to technique of analysis of variance for split- plot design by " GENSTAT Version 12th 2009 " computer software package (VSN International, Hemel Hempstead, UK) to determine The differences among treatment means were compared by L.S.D. test at $P \leq 0.05$ Payne, *et al* (2009). Combined analysis for the two seasons of experimentation was done according to the homogeneity of experimental error variance (Bartlett, 1937).

Extraction and estimation of Stevia sweeteners in relation to total**Sweeteners****. Preparation of leaves for extraction**

Stevia leaves were dried in an electric oven (E. Schulz & Co. Inh. Franz. Skorzewsh KG) at 50°C.

2. Stevia sweeteners

Stevia sweeteners were obtained from Stevia International Company for Agra industrial Projects (SKAP)

Stevioside standard preparation

Stevioside standard preparation was carried out according to Nishiyama *et al.* (1992) as follows: Dried Leaves (10 g leaves of *Stevia rebaudiana*,

Bertoni obtained from Sigma) were extracted by soaking leaves in 1.0 liter of water nearly boiling (85°C) for 30 minutes. The resulting liquid fraction was separated by Buchner filtration and the residue was washed with an additional volume of hot water (50 ml). The aqueous solution was concentrated by lyophilization (Edwards Model EF03, England) to be 50 ml and defatted by ethyl acetate, then extraction with isobutyl alcohol (150 ml) was carried out. The aqueous phase was discarded and the organic solution was evaporated by rotary evaporator (Type 349, James Jobling and Co. Ltd., England) at 70°C until drying was obtained. The dried extract was dissolved in hot methanol (100 ml) and kept over night to crystallize. The crystals were separated by filtration and re-dissolved again in boiling methanol (50 ml). This solution was clarified with active charcoal (B.D.H. Laboratory Chemicals Division. Poole, England) and left to recrystallize. The procedure was repeated three times until the formation of colorless crystals was observed. The pure solution of the stevioside prepared (pH of 9) was subjected to HPLC. At the same time an authentic pure stevioside was obtained from Nu Naturals Inc. U.S.A.

. Extraction of Stevia sweeteners from Stevia leaves with

methanol (MeoH)

Extraction of Stevia sweeteners from leaves were carried out by 0.5gm of dry stevia leaves was ground and dissolved in 0.5 ml methanol and put in shaking and heating for 30 minutes at 70°C then kept in room temperature for cooling then abukhner funnel was used for filtration using A filter paper one time after

that we used Activated charcoal for filtration another time finally we kept the filtrate frozen until analysis

Identification and quantitative analysis of Stevia sweeteners by HPLC in Stevia leaves extract

Stevia (*Stevia rebaudiana* Bertoni) leaves extract was separated and identified on HPLC as follows: Stevioside and other sweet components standard as prepared above were filtered through a millipore membrane (13 mm. diameter, 0.5 μ m pore size) were subjected for determination by chromatography with stevioside standard as internal standard. Different extracts of Stevia leaves were injected for chromatography Acetonitrile with HPLC grade (Fisons Co. England) was used in this study as mobile phase. HPLC separation was carried out on 210nm (Agilent 1200PDA detector); Eclipse plus C18 column (3.5 μ m 4.6x250 mm); linear gradient over 20 min (84:55% CH₃CN in H₂O/ 0.1% TFA); flow rate 2.0 mL/min. Injection volume: 70 μ l at ambient temperature (25°C). All the conditions used were according to **Makapugay et al., (1984)**. For each sample identification, quantification and the retention time were as described by **Makapugay et al. (1984)**. Area under each peak was used to calculate the percent of each compound.

Results and Discussion

Effect of nitrogen fertilizer levels

Results in Table (1) indicate that dry leaves yield (kg/fed), Stevioside yield (kg/ fed / year) and rebaudioside A yield (kg / fed / year) were significantly affected by increasing nitrogen fertilizer levels in the first and second seasons. The highest nitrogen rate of 40 kg N /fed/ cut gave the highest values as compared with the other nitrogen treatments of all studied characters in both seasons. The highest values of dry leaves yield were (3640 and 4025kg), respectively, in 2015/2016 and 2016/2017 seasons. These results are in agreement with **Attia (2005)**, **Lee et al. (1980)**, **Chalapahti et al. (1999)** and **Ângelo et al (2017)**. Plants fertilized with 40 N kg / fed / cut had the highest values of stevioside%. recorded (6.09 and 6.62%) respectively, in 2015/2016 and 2016/2017 seasons. Similar results were obtained by **Attia (2005)**. Adding 40 kg. N/fed/cut increased the values of rebaudioside A% (7.23 and 8.04%) in 2015/2016 and 2016/2017 season, respectively. Similar results were obtained by **Geeta and Midmore (2017)**. Nitrogen fertilizer levels of 40 kg .N/fed/ cut gave the highest values of Stevioside yield (kg /fed/ year) (238.9 and 286.1 kg) respectively, in the first and second seasons. Plants fertilized with 40 N kg / fed / cut had the highest values of rebaudioside A (kg /fed/year) (287.2 and 350.5 kg)) respectively, in the first and second seasons. On the other hand, the lowest nitrogen rate of 20 kg N/fed cut produced the lowest values of all studied characters in both seasons.

Discussion Nitrogen levels from 20 to 30 and 40 kg

.N/fed/ cut these increases may be attributed to the increase in content percentage Steviol Glycoside (SG), rebaudioside- A, in the leaves. These results are in agreement with those obtained by **Rodrigues et al (2017)**, **Attia (2005)**. **Geeta and Midmore (2017)**.

Effect of cultivars

Data presented in Table (2) indicate clearly that stevia varieties were significantly different in dry leaves yield (kg/fed), Stevioside yield (kg / fed / year) and rebaudioside A yield (kg / fed / year) in the first and second seasons. China1 varieties gave the highest values of all studied characters except dry leaves yield kg/fed in the first and second seasons.

Egy1 Varieties gave the highest values dry leaves yield in the first and second seasons respectively. China1 varieties produced the highest values of stevioside%, respectively in 2015/2016 and 2016/2017 seasons. Similar results were obtained by **Asmaa Haraz (2016)**. The highest values of rebaudioside A% were produced from china1 varieties in the first season. The current work is in agreement with **Asmaa Haraz (2016)** and **Anami et al., (2016)**. China 1 varieties produced the highest values of stevioside kg /fed yield/ year, in 2015/2016 and 2016/2017 season. Similar results were obtained by **Thiyagarajan and Venkatachalam (2015)** and **Anami et al., (2010)**. The highest values of rebaudioside A (kg /fed/ year) was produced from china1 varieties respectively in the first seasons. The current work is in agreement with **Asmaa Haraz (2016)**. Mean while, spanti varieties gave the lowest values of all studied characters in both seasons.

Discussion

Such variation might reflect the efficiency of plant building metabolites or might be ascribed to genetical differences. These results are in agreement with those found by **Asmaa Haraz(2016)** and **Thiyagarajan and Venkatachalam (2015)**

Effect of interaction between nitrogen fertilizer levels and stevia cultivars

Data in Table (3) show that the interaction effect between nitrogen fertilizer levels and stevia varieties were significant for dry leaves yield kg/fed, Stevioside yield (kg / fed / year) and rebaudioside A yield (kg / fed / year) . Nitrogen fertilizer levels of 40 kg N/ fed/cut and China1 varieties gave the highest values of all studied characters except dry leaves yield in the first and second seasons. Nitrogen fertilizer level of 40 kg N/ fed/cut and egy1 varieties gave highest values of dry leaves yield kg/fed in 2015/2016 and 2016/2017 season. Adding 40kg. N /fed / cut and China1 varieties recorded the highest values of stevioside% in 2015/2016 and 2016/2017 seasons. The highest values of rebaudioside A% were obtained by treating China1 varieties with nitrogen rates of 40 kg N /fed / cut , respectively in the first and second seasons.

Table 1. Effect of nitrogen fertilizer levels on yield and chemical composition of stevia in 2015/2016 and 2016/2017 seasons.

characters	Dry leaves yield kg /fed/ year		Stevioside%		Rebaudioside A%		Stevioside kg /fed/ year		Rebaudioside A kg /fed/ year	
	2015/2016	2016/2017	2015/2016	2016/2017	2015/2016	2016/2017	2015/2016	2016/2017	2015/2016	2016/2017
N1	2088	2651.	2.98	3.28	3.69	3.93	69.6	99.4	87.6	119.3
N2	2743	3088.	4.59	4.91	5.44	6.01	135.3	162.2	159.7	199.3
N3	3640	4025	6.09	6.62	7.23	8.04	238.9	286.1	287.2	350.5
Mean	2824	3255	4.55	4.93	5.45	5.99	148.0	182.6	178.2	223.0
L s d 0.05	262.2	253.8	—	—	—	—	22.03	9.57	28.15	11.44

N1: 20 kg N fed/cut. , **N2:** 30 kg N fed/cut , **N3:** 40 kg N fed/cut.

Table 2. Effect of cultivars on yield and chemical composition of stevia in 2015/2016 and 2016/2017 seasons.

characters	Dry leaves yield		Stevioside%		Rebaudioside A%		Stevioside kg /fed yield/ year		Rebaudioside A kg /fed yield/ year	
	2015/2016	2016/2017	2015/2016	2016/2017	2015/2016	2016/2017	2015/2016	2016/2017	2015/2016	2016/2017
Treatment										
V1	3355	3717	6.83	7.34	8.44	9.26	242.7	286.1	300.8	363.2
V2	3894	4442	5.00	5.52	6.04	6.63	204.4	256.3	245.3	307.5
V3	1729	1889	2.07	2.32	2.22	2.61	39.7	50.2	41.7	56.0
V4	2316	2970.	4.31	4.56	5.12	5.47	105.0	137.8	124.9	165.5
Mean	2824.	3255.	4.55	4.93	5.45	5.99	148.0	182.6	178.2	223.0
L s d 0.05	340.8	294.1	—	—	—	—	20.25	13.74	24.85	16.59

V1: China 1 **V2:** Egy 1 **V3:** Spanti , **V4** : Shou 2

Table 3. The interaction effect of between nitrogen fertilizer levels and stevia cultivars on chemical composition of stevia in 2015/2016 and 2016/2017 seasons

Character s	Dry leaves yield		Stevioside%		Rebaudioside A%		Stevioside kg /fed yield/ year		Rebaudioside A kg /fed yield/ year	
	2015/2016	2016/2017	2015/2016	2016/2017	2015/2016	2016/2017	2015/2016	2016/2017	2015/2016	2016/2017
Treatment										
N1V1	2222	2701	4.89	5.22	5.87	6.26	108.7	141.0	130.3	169.2
N1V2	3009	3693	3.79	4.32	5.15	5.18	114.0	159.6	154.9	191.5
N1V3	1346	1351	0.34	0.33	0.33	0.41	4.6	4.5	4.4	5.4
N1V4	1773.	2860	2.89	3.23	3.42	3.88	51.2	92.5	60.6	111.0
N2V1	3380	3735	7.06	7.61	8.93	9.63	238.7	284.1	301.8	359.6
N2V2	3848.	4155	4.28	4.66	4.52	5.59	164.8	193.7	174.0	232.4
N2V3	1443	1764.	2.76	2.95	3.26	3.54	98.0	52.0	47.1	62.4
N2V4	2302	2698	4.26	4.42	5.04	5.3	380.6	119.1	116.1	143.0
N3V1	4463	4716	8.53	9.18	10.54	11.89	380.6	433.1	470.2	560.8
N3V2	4824	5477	6.93	7.59	8.44	9.10	334.5	415.5	407.0	498.6
N3V3	2399	2552.	3.11	3.69	3.07	3.92	74.7	94.2	73.7	100.1
N3V4	2875.	3353.	5.77	6.02	6.88	7.23	165.8	201.8	197.9	242.5
mean	2824.	3255.	4.55	4.93	5.45	5.99	148.0	182.6	178.2	223.0
L s d 0.05	541.8	475.1	—	—	—	—	34.24	21.60	42.45	26.07

N1: 20 kg N fed/cut., **N2:** 30 kg N fed/cut , **N3:** 40 kg N fed/cut.
V1: China 1 **V2:** Egy 1 **V3:** Spanti , **V4** :Shou 2

Nitrogen fertilizer level of 40 kg N/ fed/cut and china 1 varieties gave highest values of stevioside (kg /fed / year) in 2015/2016 and 2016/2017 seasons, respectively. Adding 40kg. N /fed / cut to Chinal varieties recorded the highest values of rebaudioside A (kg /fed/year) respectively in the first and second seasons.

References

- Attia.E. A (2005)** Effect of some agronomic treatments on stevia (*stevia rebaudiana*,bertoni) yield and quality in egypt. Ph D thesis.Fac of AgricAin shams Univ. Egypt.
- Asmaa Haraz S, M (2016)** Genetic improvement through selection of Stevia rebaudiana under Egyptian conditions. M.Sc.thesis, Fac of Agric Saba Basha . Alexandria Univ. Egypt.
- Anami, E. T, J. P. Poletine, M. C. Gonçalves-Vidigal, P. S. V. Filho, G. F. Lacanallo, M. V. Kvitschal and A. Gonela (2010).** Characterization and genetic divergence in *Stevia rebaudiana*(Bert.)Bertoni clones based in agronomical and morphological characteristics. Journal of Food, Agriculture & Environment.8 (3&4) : 4 6 3 - 4 6 9
- Bartlett, M. S. (1937).** Properties of sufficiency and statistical tests. Proceedings of the London. Series A-Math. Phys. Royal Society of Sci., 160(901): 268 282.
- Chalapathi, M. V., S.Thimmegowda, G. G. E.Rao, N. Devakumar and J.Chandraprakash (1999).**Influence of fertilizer level on growth, yield and nutrient uptake of ratooncrop of stevia (*Stevia rebaudiana*). Journal of Medical and Aromatic Plant Sci.21 : 947-949.
- Kovacevi. D. B, F. J. Barba, D. Granato, C. M.Galanakis, Z. Herceg, V. D. Uzelac and P.Putnik (2018)** Pressurized hot water extraction (PHWE) for the green recovery of bioactive compounds and steviol glycosides from *Stevia rebaudiana* Bertoni leaves. Food Chemistry. Vol 254, 15 July, 150-157
- Putnik Geuns, J. M. C. (2003).**Stevioside.Phytochemistry, 64(5): 913–921.
- Geeta. G.K, Dd. J. Midmore and R.Gautam(2017)** Effect of nutrient omission and pH on the biomass and concentration and content of steviol glycosides in stevia (*Stevia rebaudiana* (Bertoni) Bertoni) under hydroponic conditions. Journal of Applied Research on Medicinal and Aromatic Plants.Vol 7, December , 136-142
- Lee, J. I, K. H. Kang, H. W. Park,Y. S Ham and C. H Park(1980)** Studies on the new sweetening source plant, Stevia rebaudiana in Korea. II. Effects of fertilizer rates and planting density on dry leaf yields and various agronomic characteristics of Stevia rebaudiana. Research Reports of the Office of Rural Development, Crop, Suwon Vol.22: 138-144
- Makapugay, H. C., N.P.D. Nanayakkara and A.D. Kinghorn (1984).** Improve high-performance liquid chromatographic separation of the *Stevia rebaudiana*sweetditerpene glycosides using linear gradient elution. Journal of Chromatography A.283 : 390-395
- Negar T., Y. Hamidoghli, B. Rabiei and S. Hamidoghli (2012)**Effects of plant density and transplanting date on herbage, stevioside, phenol and flavonoid yield of Stevia rebaudiana Bertoni. Intl J Agra Crop Sci. Vol., 4 (6), 298-302
- Nishyama, P., M. Alvarez and L. G. E. Vieira (1992).**Quantitative analysis of stevioside in the leaves of *Stevia rebaudiana*bynea r infrared reflectance spectroscopy. Journal of Science Food and Agriculture. 59(3) 277-281.
- Megeji . N. W., J. K. Kumar, Virendra Singh, V. K. Kaul and P. S. Ahuja 2005** Introducing Stevia rebaudiana, a natural zero-calorie sweetener. Current Sci Vol.(5) : 801-804
- Payne, R. W., D. A. Murray, S. A. Harding, D. B. Baird and D. M. Soutar (2009).** Gen Stat for Windows (12th Edition) Introduction. VSN International, Hempstead, UK.
- Rodrigues. A .M , SandraAfonso, Isabel Q. Ferreira and Margarida. A (2017)** Response of stevia to nitrogen fertilization and harvesting reime in northeastern Portugal .journl Archives of Agronomy and Soil Science . Volume 63 : 626-637
- Roberto .L.M, A. V.Gálvez, Liliana Z..B and Kong A .H (2012)** Stevia rebaudianaBertoni, source of a high-potency natural sweetener: A comprehensive review on the biochemical, nutritional and functional aspects . Food Chemistry 132 :1121–113
- Singh. B, J. Singh, A. Kaur (2014)**Agro-production, Processing and Utilization of *Stevia rebaudiana*as Natural Sweetener . Journal of Agri Engineering and Food Technology; Vol 1, (1) 28-31
- Soejarto .D.D, Compadre .C.M, Medon PJ, Kameth .S.K, Kinghorn .A.D(1983).** Potential sweetening agents of plant origin. II. Field search for sweet tasting *Stevia* species. Econ Bot; 37: 71-9
- Thiyagarajan, M and P. Venkatachalam (2015).**Assessment of genetic and biochemical diversity of*Steviarebaudiana*Bertoni by DNA fingerprinting and HPLC analysis.Journal of Annals of Phytomedicine. 4(1): 79-85.

التقييم الكمي لبعض أصناف الاستيفيا تحت مستويات مختلفة من التسميد الأزوتي

ابوبكر عبد الوهاب طنطاوي* احمد السيد عطية** ناريمان نايف قناوي**

*كلية الزراعة جامعة المنيا**مركز البحوث الزراعية - الجيزة

اجريت الدراسة في محطة بحوث الجيزة مركز البحوث الزراعية موسمي 2015-2016 و 2016-2017 لدراسة تأثير التسميد النتروجيني (20 ، 30 و 40 كجم نتروجين للحشة) وبعض الاصناف

(spantil, china 1, egypt 1 و shou 2) وتفاعلهما علي الصفات تحت الدراسة في الموسمين.

* اعطت معدلات التسميد الأزوتي اختلافات عالية المعنوية لجميع الصفات تحت الدراسة عند مستوي معنويه 5 % و اعطي معدل تسميد 40 كجم نتروجين/ فدان / حشة اعلي القيم لجميع الصفات محتوى الاوراق من المحليات مثل (ال Stevioside و rebaudioside-A) وكذلك محصول الفدان من المحليات مثل (ال Stevioside و rebaudioside-A) في السنه مقارنة بالمعاملات الاخرى في الموسمين وكذلك المحصول الجاف الاوراق .

*أوضحت النتائج وجود فرق معنوي كبيرة بين جميع الاصناف عند مستوي معنويه 5 % وسجل china 1 أعلى تركيز من ال-rebaudioside (A) و أعلى تركيز من ال (Stevioside) وكذلك محصول الفدان من المحليات مثل (ال Stevioside و rebaudioside-A) في السنه في الموسمين مقارنة بالاصناف الاخرى

* أوضحت النتائج أيضا ان التفاعل بين اضافة معدل 40 كجم نتروجين/ فدان / حشة والصنف china 1 اعطي اعلي القيم لجميع الصفات ما عدا كميته محصول الفدان من الاوراق الجافه حيث سجل التفاعل بين اضافة معدل 40 كجم نتروجين للحشة والصنف egypt 1 اعلي كميته محصول الفدان من الاوراق الجافه في الموسمين .